

THE SOUTHERN VILNIUS PHOTOMETRIC SYSTEM. IV. THE E REGION STANDARD STARS

M. C. Forbes¹, R. J. Dodd² and D. J. Sullivan¹

¹*Victoria University of Wellington, P.O.Box 600, Wellington, New Zealand*

²*Carter Observatory, P.O. Box 2909, Wellington, New Zealand*

1. INTRODUCTION

In the paper by Forbes, Dodd & Sullivan (1993), hereafter referred to as Paper I, measurements of the equatorial stars, already observed in the northern hemisphere Vilnius system, were made using Carter Observatory's set of filters. It was concluded that these could be accurately transformed to the standard Vilnius system and so a program to establish a set of Vilnius standard stars in the southern hemisphere was begun.

The first results of this program are given by Forbes, Dodd & Sullivan (1994), hereafter Paper II, listing a set of bright standard stars spread across the southern sky. To extend the standards to fainter stars it was decided to measure the Harvard E Region stars, which have already been established as standards in other photometric systems (e.g. Menzies et al. 1989). The Harvard E Regions consist of nine groups which contain approximately sixty stars each, ranging from $V = 3$ to 12, centred near $\delta \approx -45^\circ$ and spaced 2 – 3 hours apart in right ascension. Some measurements of the Harvard E and F Region stars are given by Zdanavičius, Dodd, Forbes & Sullivan (1995), hereafter Paper III.

2. OBSERVATIONS AND REDUCTIONS

The observations were made at the Mt. John University Observatory (University of Canterbury), situated near Lake Tekapo in the South Island of New Zealand, over the period 1992 to 1993. A set of stars previously measured in the Vilnius system (North 1980), which are observable from the southern hemisphere, were selected

as primary standards, to enable transformation of the local Vilnius filter system to the standard system. Measurements of the primary standard stars were interspersed with those of the southern standard stars. Certain southern standard stars were used as extinction stars. These were selected from F5–G2 V spectral type stars with $\delta \approx -45^\circ$ and measured every hour, using a variation of Nikonov’s method (Nikonov 1953, Straizys 1992). More details on the observing and reduction procedures are given in Paper I.

To date, 96 stars have been observed as possible standard stars. The following criteria were applied to select the stars that have been measured to acceptable precision and accuracy. First, the star must have been measured on at least three different nights, enabling a mean value and standard deviation to be calculated. Next, the standard deviation of the mean for the V magnitude and colors of the star must be less than 0.02 mag. The only exception was the color $U - P$, whose rejection threshold was relaxed to 0.025 mag. The slightly larger errors in the U filter observations imply that longer integration times are required for measurements using that filter. Finally, using either the ‘stellar box’ or $Q(\lambda)$ pseudo-spectrum methods (Straizys 1992), the measurements must enable a star to be classified within two sub-classes and within one class of its MK spectral and luminosity classes, respectively. This left 60 stars suitable for use as standards, with most of the rejections due to an insufficient number of observations per star. The rejected stars remain in the observing program, to see if further observations will enable them to meet the standard star criteria.

3. CATALOG

Table 1 lists the observations of the 60 E region standard stars measured in the southern Vilnius photometric system. The first column is the star name as given by Menzies (1989). The next two columns are the Epoch 2000 equatorial coordinates. These are followed by the Vilnius V magnitude and the six color indices. A colon following the measurement indicates that the standard deviation of the mean exceeds 0.015 mag; a double colon means the standard deviation is greater than 0.02 mag. The number of observations (n) of each star is given in the next column. The final two columns give the spectral type as listed by Menzies et al. (1989) and classified from the photometric color indices by K. Zdanavičius. Close

binaries were not classified photometrically since their color indices belong to common light of the two components.

4. DISCUSSION

The overall accuracy and the precision of the observations can be seen in Table 2. This shows the standard deviation ε of each color calculated using

$$\varepsilon = \frac{\sigma}{\sqrt{\frac{\sum n}{N}}}$$

where σ is the typical standard deviation of a single observation for the southern system, n is the number of observations of a star and N is the number of observed stars. The accuracy required for establishing a set of standard stars, i.e. 0.01 mag, has clearly been achieved.

ACKNOWLEDGMENTS. The authors thank University of Canterbury and Mt. John University Observatory staff for the generous use of their facilities, the Vilnius Institute of Theoretical Physics and Astronomy for manufacturing the filter sets and the New Zealand Lottery Board for financing their purchase. The New Zealand Foundation for Research, Science and Technology and the Internal Research Grant Committee of Victoria University of Wellington provided partial funding of this project. It is a pleasure to thank K. Zdanavičius for photometric classification of the observed stars and V. Straižys for his help and encouragement in establishing this program.

Table 1. Catalog of standard stars in the southern Vilnius photometric system (E regions)

Star	$\alpha(2000)$	$\delta(2000)$	V	$U-P$	$P-X$	$X-Y$	$Y-Z$	$Z-V$	$V-S$	n	Sp	
E136	01 ^h 15 ^m 07 ^s	-45°32'	5.05	0.49	0.61	0.76	0.32	0.20	0.51	43	G0 V	F9.5 IV
E345*	06 25 43	-48 11	5.78:0.55::0.68	0.23	0.10	0.04	0.12	5	B9 V			
E347	06 49 54	-46 37	5.12:0.47::0.58	0.63	0.29	0.18	0.45	4	F5 V		F5 V	
E428	09 30 33	-45 34	6.60	0.64::1.16	1.54	0.49	0.39	0.82	8	K2-3 III	K2 III	
E429	09 20 15	-45 10	6.67	0.66::1.21	1.61	0.49	0.41	0.84	8	K2 III	K2 III	
E439	09 12 14	-45 50	6.66	0.56	0.59	0.60	0.28	0.17	0.45	32	F3-5 V	F3 V
E441*	09 15 14	-45 33	6.27	0.49: 0.61	0.20	0.10	0.05	0.15	8	B8 V		
E444	09 22 24	-46 03	5.74	0.57: 0.89	1.14	0.44	0.27	0.67	8	G6-8 III	G7 III	
E445	09 29 22	-43 11	6.61	0.54: 0.62	0.66	0.29	0.17	0.46	8	F5 IV	F6 IV	
E446	09 31 19	-47 57	6.56	0.53: 0.73	0.50	0.20	0.12	0.30	8	A9 IV-V	A9 V	
E447	09 38 01	-43 11	5.51	0.54: 0.88	1.25	0.47	0.29	0.69	8	G8 II	G8.5 III	
E478	09 14 08	-44 09	5.88	0.27: 0.45	0.17	0.08	0.03	0.10	8	B5 V	B6 V	
E479*	09 14 24	-43 14	5.27	0.25:0.39	0.16	0.07	0.03	0.10	8	B3-5 V		
E493	09 11 33	-46 35	5.82	0.10: 0.19	0.11	0.05	0.01	0.06	8	B2 III-IV	B1.5 V	
E494	09 13 34	-47 20	5.95	0.55::0.73	0.25	0.10	0.04	0.14	6	B9 V	B9.5 V	
E496	09 16 58	-45 01	6.78	0.57::0.81	0.29	0.11	0.05	0.13	6	A0 V	A0.5 V	
E497	09 23 25	-48 17	6.32	0.27::0.35	0.16	0.08	0.03	0.11	6	B3 IV	B4 IV	
E498	09 23 39	-46 55	6.26	0.43::0.51	0.19	0.09	0.04	0.13	6	B7 III	B7 V	
E540	12 03 38	-42 26	5.17	0.52	0.58	0.60	0.27	0.16	0.43	52	F5 V	F4 V
E612	14 56 14	-44 05	7.92	0.61	0.83	0.45	0.17	0.10	0.21	4	A3 IV	A5 V
E613	14 55 25	-44 20	8.02	0.67	0.81	0.49	0.19	0.12	0.26	4	A4 V	A5.5 IV
E614	14 53 04	-43 44	8.16	0.66	0.81	0.49	0.20	0.13	0.29	4	A5 V	A6.5 IV
E633	14 38 12	-45 52	6.90	0.64::1.18	1.52	0.48	0.38	0.82	4	K2 III	K2 III	
E639*	14 30 08	-45 19	5.51	0.53	0.60	0.20	0.10	0.04	0.12	15	B8 V	
E640	14 36 18	-46 15	5.58	0.76	1.34	1.93	0.59	0.49	0.99	14	K3 III	K3.5 III
E641	14 39 11	-46 35	6.11	0.51	0.58	0.69	0.30	0.19	0.48	7	F6 V	F7 V
E642*	14 45 57	-44 52	6.95	0.56	0.79	0.32	0.14	0.07	0.15	5	A0 V	A0 V
E643	14 46 28	-47 26	5.77	0.61	0.83	0.36	0.14	0.06	0.16	7	A1 V	A1.5 V
E644	14 47 31	-43 33	6.34	0.63	1.00	1.32	0.49	0.32	0.76	17	G8 III	G9.5 III

Table 1 (continued)

Star	$\alpha(2000)$	$\delta(2000)$	V	$U-P$	$P-X$	$X-Y$	$Y-Z$	$Z-V$	$V-S$	n	Sp	
E645	14 ^h 56 ^m 24 ^s	-44°42'	6.77	0.62	0.83	0.35	0.13	0.07	0.15	35	A1 V	A1.5 V
E646*	14 59 18	-43 28	7.08	0.56	0.66	0.57	0.24	0.16	0.38	4	F2 III	
E647	14 59 45	-43 49	6.60	0.49	0.55	0.66	0.29	0.18	0.46	65	F7 IV-V	F5 V
E679*	14 26 10	-45 23	4.38	0.72	0.81	0.57	0.27	0.17	0.46	9	F7	
E680	14 27 12	-46 08	5.85	0.62	0.76	0.58	0.21	0.14	0.32	9	A1m	A6 V
E681*	14 28 51	-47 59	6.41	0.43	0.53	0.19	0.08	0.05	0.11	8	ApSi	
E682	14 35 09	-46 28	6.92	0.68	0.68	0.23	0.13	0.06	0.16	8	B9 V	B9 III
E684	14 35 49	-46 49	6.82	0.61	0.89	1.18	0.46	0.29	0.71	8	G8 III	G7 III
E687	14 37 54	-48 01	6.66	0.53	0.58	0.22	0.13	0.06	0.17	4	B8 V	B8 III-IV
E688*	14 40 18	-45 48	6.64	0.42:	0.51	0.20	0.04	0.07	0.15	3	ApSi	
E690	14 50 58	-42 49	6.84	0.65	0.84	0.43	0.15	0.09	0.21	4	A2 V	A4 V
E691*	14 51 37	-43 34	4.34	0.28	0.33	0.15	0.07	0.04	0.10	8	B5 IV	
E695	14 59 26	-43 10	6.14	0.74	0.72	0.78	0.32	0.22	0.54	6	F7 II	F7 II
E6100	14 54 00	-43 36	8.23	0.59:	0.91	1.30	0.48	0.31	0.73	4	G8 II	G9 III
E748	17 24 43	-45 01	6.70	0.54	0.60	0.58	0.26	0.15	0.40	95	F3 V	F2.5 V
E765	17 37 53	-42 34	7.22	0.48	0.65	0.83	0.34	0.22	0.56	55	G2-3 V	G2 IV-V
E869	19 55 15	-41 52	4.14:	0.62	1.00	1.35	0.47	0.34	0.76	4	K0 II-III	K0 III
E924	22 32 11	-43 16	6.99	0.59	0.95	1.23	0.45	0.29	0.72	6	K0 III	G9.5 III
E929	22 33 54	-44 41	6.87	0.58	0.87	1.16	0.45	0.28	0.71	7	G8 III	G6 III
E930	22 53 15	-45 09	6.93	0.63	1.11	1.42	0.49	0.34	0.78	7	K0 III	K0.5 III
E940	22 30 43	-44 06	7.00	0.62	0.84	0.37	0.14	0.07	0.15	5	A0-1 V	A2 V
E942	22 36 41	-43 28	6.81	0.52	0.57	0.70	0.31	0.20	0.49	37	F8 V	F6 V
E943	22 42 43	-44 15	6.15	0.58	0.90	1.19	0.45	0.29	0.71	7	K0 III	G8 III
E944	22 45 41	-46 33	5.60	0.69	1.23	1.71	0.55	0.41	0.89	7	K2 III	K2.5 III
E948	22 30 15	-42 17	6.99	0.57	0.74	0.25	0.11	0.04	0.11	7	B9.5 V	B9.5 V
E949	22 47 09	-41 42	6.92	0.67	1.22	1.66	0.53	0.41	0.88	6	K2 III	K2.5 III
E963*	22 42 37	-47 12	6.05	0.49	0.59	0.74	0.32	0.20	0.53	4	G0 V	

Table 1 (continued)

Star	$\alpha(2000)$	$\delta(2000)$	V	$U-P$	$P-X$	$X-Y$	$Y-Z$	$Z-V$	$V-S$	n	Sp
E971	22 ^O 15 ^m 35 ^s	-44 ^O 27'	6.18	0.62	0.94	1.26	0.48	0.30	0.73	6	G8-K0 III G8.5 III
E972	22 23 07	-45 56	5.69	0.56	0.63	0.57	0.25	0.15	0.40	6	F0 V F1.5 V
E973	22 23 25	-46 40	6.96	0.65	0.83	0.36	0.13	0.08	0.16	4	A1 V A1.5 V
E979	22 43 30	-41 25	4.91	0.61	0.97	1.26	0.47	0.31	0.74	5	K0 III G9.5 III

N o t e s (from Menzies 1989):

E345 – double, 6.0 + 8.3, 1.5''

E441 – double, 6.6 + 7.7, 1''

E479 – double, 5.2 + 9.6, 6''

E639 – companion approx 8'' NW

E642 – double, 7.3 + 12.0, 35'', companion not included

E646 – double, 7.2 + 11.2, 4''

E679 – double, 5.2 + 5.3, 0.3'', cpm

E681 – companion 8'' NW

E688 – companion approx 8'' W

E691 – double, 0.1''

E963 – double, 6.4 + 9.8, 8'', cpm

Table 2. Summary of precision of the Vilnius southern standard stars.

V	$U-P$	$P-X$	$X-Y$	$Y-Z$	$Z-V$	$V-S$
0.008	0.008	0.005	0.004	0.004	0.004	0.005

REFERENCES

- Forbes M. C., Dodd R. J., Sullivan D. J. 1993, *Baltic Astronomy*, 2, 246
Forbes M. C., Dodd R. J., Sullivan D. J. 1994, *Baltic Astronomy*, 3, 227
Zdanavičius K., Dodd R. J., Forbes M. C., Sullivan D. J. 1995, *Baltic Astronomy*, 4, 25
Menzies J. W., Cousins A. W. J., Banfield R. M., Laing J. D. 1989, *Circ. South African Astron. Obs.*, No. 13, 1
Nikonov V. B. 1953, *Bull. Abastumani Obs.*, No. 14
North P. 1980, *A&AS*, 41, 395
Straižys V. 1992, *Multicolor Stellar Photometry*, Pachart Publishing House, Tucson, Arizona